

TTP-04-11
 WUE-ITP-2004-015
 August 2004

TESTS OF THE NONCOMMUTATIVE STANDARD MODEL AT A FUTURE PHOTON COLLIDER^a

THORSTEN OHL^{1b} and JÜRGEN REUTER^{2cd}

¹*Institut für Theoretische Physik und Astrophysik,
 Universität Würzburg, D-97074 Würzburg, Germany*

²*Institut für Theoretische Teilchenphysik,
 Universität Karlsruhe, D-76128 Karlsruhe, Germany*

Extensions of the Standard Model of elementary particle physics to noncommutative geometries are proposed by string models. Independent of this motivation, one may consider such a model as an effective field theory with higher-dimensional operators containing an antisymmetric rank-2 background field. We study the signals of such a Noncommutative Standard Model (NCSM) and analyze the discovery potential of fermion pair production at a future photon collider.

The interest in noncommutative (NC) theories has been growing dramatically in recent years due to the observation that open string theories with constant antisymmetric rank-2 tensor fields can be interpreted as Yang-Mills theories living on a NC manifold¹. Independent from this motivation, NC Quantum Field Theory in itself provides an interesting approach in introducing a fundamental length scale and cutting off short-distance contributions, consistent with the symmetries of a given model. Recently, there has been a lot of effort in constructing an Effective Field Theory (EFT) which is defined on a NC spacetime with a canonical structure

$$[\hat{x}^\mu, \hat{x}^\nu] = i\theta^{\mu\nu} = i\frac{1}{\Lambda_{\text{NC}}^2}C^{\mu\nu} = i\frac{1}{\Lambda_{\text{NC}}^2}\begin{pmatrix} 0 & -E^1 & -E^2 & -E^3 \\ E^1 & 0 & -B^3 & B^2 \\ E^2 & B^3 & 0 & -B^1 \\ E^3 & -B^2 & B^1 & 0 \end{pmatrix} \quad (1)$$

and which has the Standard Model (SM) as low energy limit for $\sqrt{s} \ll \Lambda_{\text{NC}}$. The noncommutativity $\theta^{\mu\nu}$ is an antisymmetric real matrix, assumed here to be constant and understood as a spurion, breaking Lorentz invariance. Its origin is irrelevant as long as we are merely studying the EFT, where it appears

^aWork supported by Deutsche Forschungsgemeinschaft (RU 311/1-1), and Bundesministerium für Bildung und Forschung, Germany (05HT1RDA/6).

^be-mail: ohl@physik.uni-wuerzburg.de

^ce-mail: reuter@particle.uni-karlsruhe.de

^dPresented at LCWS2004, Paris, France, April 2004

as a coefficient in front of operators of dimension six or higher. The crucial theoretical point is the realization of gauge invariance on NC spaces^{1,2}. A general way is provided by the Seiberg-Witten Map (SWM)¹. It is an asymptotic expansion in the noncommutativity $\theta^{\mu\nu}$, which expresses the NC matter and gauge fields as functions of commutative matter and gauge fields so that the NC gauge transformations are realized by the commutative ones.

In the Noncommutative Standard Model (NCSM)³, one has to replace each SM field by the corresponding SWM and all products by $\star$ -products (which realizes products of noncommutative functions in terms of commutative ones). There are higher-dimensional operators, whose contributions are suppressed by the ratios $\Lambda_{EW}^2/\Lambda_{NC}^2$ or s/Λ_{NC}^2 of the EW scale or the CMS energy and the NC scale, giving rise to deviations of SM decay rates and production cross sections. SM vertices receive corrections with new Lorentz structures, and there will be contact terms of the form $f\bar{f}\gamma\gamma$ etc., which are required by gauge invariance. In addition there can also be triple neutral gauge boson couplings (TGC) like $\gamma\gamma\gamma$, $Z\gamma\gamma$, etc.⁴. Current bounds (mainly from the non-observation of the Lorentz-violating Z decays $Z \rightarrow \gamma\gamma, gg$ at LEP) are $\Lambda_{NC} \gtrsim 100$ GeV. In a low-lying string scenario one could expect values for such a scale as low as $\Lambda_{NC} \gtrsim 1$ TeV.

In the NCSM, amplitudes are asymptotic expansions in $\theta^{\mu\nu}$. In lowest order in θ , for processes forbidden in the SM, the square of the first order NCSM amplitude is the leading contribution, while for a nonvanishing SM amplitude the deviation of the leading order is given by the interference of the SM with the lowest order NCSM amplitude. Higher orders require the (yet unknown) second and higher orders in the SWM.

Our proposal is to study fermion pair production $\gamma\gamma \rightarrow f\bar{f}$ at a future photon collider. Such a machine is planned to be operated at a future linear e^+e^- collider with energies up to 1 TeV. Highly energetic photons are produced by Compton backscattering of laser photons off the electron beam. The photons can be delivered with a high degree of polarization, which is crucial for our considerations.

Theoretically, producing massless fermions ($\sqrt{s} \gg m_f$) from polarized photons suggests to use the very concise formalism of helicity amplitudes for evaluating the cross section. The noncommutativity $\theta^{\mu\nu}$ can elegantly be converted to a spinor expression containing only the spinor metric ϵ_{AB} and a symmetric rank-2 spinor ϕ_{AB}

$$\theta_{A\dot{A},B\dot{B}} = \theta^{\mu\nu} \bar{\sigma}_{\mu,A\dot{A}} \bar{\sigma}_{\nu,B\dot{B}} = \phi_{AB} \epsilon_{\dot{A}\dot{B}} + \bar{\phi}_{\dot{A}\dot{B}} \epsilon_{AB} \quad (2)$$

with $\phi_{AB} = (\bar{\phi}_{\dot{A}\dot{B}})^*$ and three independent complex components

$$\phi_{11} = -E_- - iB_-, \quad \phi_{12} = E_3 + iB_3 = \phi_{21}, \quad \phi_{22} = E_+ + iB_+, \quad (3)$$




Figure 1: *Left*: Azimuthal dependence of the amplitudes. The horizontal line is the SM differential cross section. *Right*: The Z resonance in the interference of SM and $\mathcal{O}(\theta)$ NCSM amplitude plotted for different values of the azimuth angle.

where $E_{\pm} = E_1 \pm iE_2$ and $B_{\pm} = B_1 \pm iB_2$. Using this decomposition, all amplitudes can be expressed as contractions of Weyl-van der Waerden spinors with ϕ and among themselves.

Due to helicity conservation in the SM, the only nonvanishing combinations for the processes $\gamma\gamma \rightarrow f\bar{f}$ with massless fermions are $(\pm, \mp) \rightarrow (\pm, \mp)$ and $(\pm, \mp) \rightarrow (\mp, \pm)$. Thus a deviation from the SM can only be seen, if the two photon helicities add up to a total of two. Therefore the photon collider has to be run in the d -wave mode ($S = 2$), while the preferred mode for Higgs physics is the s -wave ($S = 0$). Fortunately, one can easily switch between them. For both SM diagrams, there are two $\mathcal{O}(\theta)$ contributions, where one SM vertex is replaced by the corresponding NCSM vertex, while there is a new contact term as well as two new s -channel diagrams containing the TGCs. A nonvanishing interference appears only for non-zero E^1 or E^2 . The explicit expressions for the different helicity amplitudes are published elsewhere⁵.

As cross checks, we verified numerically the amplitudes calculated within two completely different formalisms – helicity amplitudes vs. Dirac spinors using the optimizing matrix element compiler `O'Mega/WHIZARD`⁶ – to agree within numerical accuracy.

In the NCSM, the Z can be produced resonantly in photon collisions. The right figure 1 shows the resonance structure around the Z pole. Unfortunately this can not be well observed at the photon collider because of poor statistics




Figure 2: Number of events per year in the two halfspheres $\phi \gtrless 0$ for $\sqrt{s} = 800$ GeV in the $S = 2$ mode. A cut has been made 0.02 radians around the beam axis.

in the photon spectrum for low energies. In figure 1 on the left, the differential cross section at $\sqrt{s} = 800$ GeV is shown plotted against the azimuth angle ϕ . From this, one sees that the integrated cross section is the same for the pure SM and the interference, as the interference goes like $\propto \sin(\phi + \phi_0)$, where ϕ_0 is a phase which depends on the spatial orientation of the noncommutativity. However, in the polarized cross section it is possible to scan over the final state fermions to look for angular dependent deviations from the SM prediction.

To get a realistic result, one has to fold the cross section with the photon spectrum produced in Compton scattering for the photon collider, for which the program *Circe 2.0* has been used⁷. In figure 2 we show the number of binned events over the invariant mass of the fermion pair, assuming an integrated luminosity of $\mathcal{L} = 2000 \text{ fb}^{-1}$. The cross section and angular variation away from the Z resonance do not depend very much on the specific value of the TGCs. A signal could always be seen if the scale Λ_{NC} is even a little bit higher than the CMS energy of the linear collider. It is not consistent to consider the amplitudes and cross section for collider energies higher than the scale Λ_{NC} , since higher orders in θ can only be neglected if $s/(\Lambda_{\text{NC}})^2 \lesssim 1$.

In conclusion, the NCSM offers a variety of new phenomena due to the presence of an antisymmetric rank-2 spurion field which breaks Lorentz invariance at a scale Λ_{NC} . This violation of angular momentum conservation is a generic signal for noncommutativity, which leads to deviations from the




Figure 3: Number of events per year in the two half-spheres $\phi \gtrless 0$ for $\sqrt{s} = 500$ GeV and $\sqrt{s} = 200$ GeV in the $S = 2$ mode.

isotropic distribution of final state particles around the beam axis. We have focused on fermion pair production at a future photon collider. Polarization is shown to be a mandatory ingredient to be sensitive for signals of noncommutative theories. In a conservative estimate, a photon collider will be sensitive to scales of the order of $\Lambda_{\text{NC}} \sim 1$ TeV, but if enough data will be available, the sensitivity may even be better by a factor three to five.

References

1. N. Seiberg, E. Witten, JHEP 9909 (1999) 032 [hep-th/9908142]
2. J. Wess, Commun. Math. Phys. **219** (2001) 247.
3. X. Calmet et al., Eur. Phys. J. **C23** (2002) 363 [hep-ph/0111115].
4. G. Duplanić, P. Schupp, and J. Trampetić, Eur. Phys. J. **C32** (2003) 141 [hep-ph/0309138] and references therein.
5. T. Ohl, J. Reuter, Phys. Rev. **D** (in print) [hep-ph/0406098].
6. T. Ohl, hep-ph/0011243; M. Moretti, T. Ohl, J. Reuter, hep-ph/0102195; <http://whizard.event-generator.org>; W. Kilian, T. Ohl, J. Reuter, hep-ph/0708.4233.
7. T. Ohl, Comput. Phys. Commun. **101** (1997) 269 [hep-ph/9607454]; T. Ohl, Circe 2.0, WUE-ITP-2002-006.